



Area Symbol: OH175, Soil Area Version: 18

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Non-Irr Class *c	Alfalfa hay	Brome-grass alfalfa	Brome-grass alfalfa hay	Corn	Corn silage	Grass legume hay	Grass legume pasture	Kentucky bluegrass	Oats	Orchardgrass	Orchardgrass alfalfa	Orchardgrass alfalfa hay	Pasture	Soybeans	Tobacco	Tomatoes	Wheat	Winter wheat	*n NCCPI Soybeans	eFOTG PI
FuA	Fulton silty clay loam, 0 to 2 percent slopes	74.11	24.7%		Illw				100		4			75	6.4	4			36		17.5		40	57	64
Lb	Latty silty clay loam	57.50	19.2%		Illw				110		4.5			82			5.2		42				46	53	81
Lc	Latty silty clay	41.63	13.9%		Illw				100		4			80			5		38				40	49	78
DeA	Del Rey silt loam, 0 to 2 percent slopes	38.36	12.8%		llw		7.5		115		4.5			69			4.5		42				48	71	76
Blg1B1	Blount silt loam, ground moraine, 2 to 4 percent slopes	25.65	8.6%		lle				137		4.6	9							44				61	47	75
Blg1A1	Blount silt loam, ground moraine, 0 to 2 percent slopes	23.45	7.8%		llw				141		4.6	9.3							45				63	51	78
Br	Bono silty clay	16.33	5.4%		Illw				115		4			78			4.3		40				40	55	76
Gwg1B2	Glynwood silt loam, ground moraine, 2 to 6 percent slopes, eroded	11.08	3.7%		lle				128		4.2	8.5							44				57	26	65
Pa	Pandora silty clay loam	4.70	1.6%		llw				120		4.5			70				8	48				48	67	71
Lk	Lindsay silt loam, occasionally flooded	2.75	0.9%		llw	4.5			115		4.5		5.5						48	2800		40		71	80
NpB	Nappanee silt loam, 2 to 6 percent slopes	2.37	0.8%		Ille	5.5		3.8	95	17	4.5			85					34				38	52	70
MrD2	Morley silt loam, 12 to 18 percent slopes, eroded	1.72	0.6%		Ive				106	15.6	3.5	7.8							37				47	30	59
PkA	Pewamo silty clay loam, 0 to 1 percent slopes, eroded	0.28	0.1%		llw				157		5	10.5							47				64	67	84
Weighted Average						0.1	1		112.6	0.2	4.3	1.9	0.1	60.2	1.6	1	2.5	0.1	40.4	25.7	4.3	0.4	46.2	*n 54.4	73.8

Soils data provided by USDA and NRCS.